

# Understanding the stability of positive electrode materials for aqueous organic redox flow batteries

Sri Narayan

Bo Yang, Advait Murali, Vinayak Krishnamurthi and G.K. Surya Prakash  
*University of Southern California, Los Angeles, CA*

Presentation ID# 405

Session Topic : Flow Batteries

DoE Office of Electricity  
Peer Review Meeting  
October 27, 2021

# Goals and Objectives

## Overall Goal



Advance the understanding of an All-Organic Water-based RFBs for cost-effective large-scale energy storage.

## Specific Objectives

- Understand the mechanisms of degradation to formulate design rules for new molecules
- Design and demonstrate inexpensive positive side organic molecules that can be cycled repeatedly without degradation


# Advances Made at USC in RFBs

- First ever aqueous all-organic redox flow batteries based on quinone derivatives (BQDS/AQDS) in acidic media.
- Mitigated crossover by molecular design and symmetrical cycling.
- Demonstrated cycling of Michael-Reaction resistant positive side materials (DHDMMBS)
- Scale up and demonstration (ITN) of 1kW/2KWh all-organic redox flow batteries (DHDMMBS/AQDS) 4000 Ah/ 4.5 h per cycle with <0.01% loss/hour.
- Developed Iron sulfate/AQDS system for stable cycling over 1000 cycles with < 8 x10<sup>-6</sup> %/hour.



# Capacity Degradation Modes and Rates

- For realizing LCOS of < 5 cents/kWh degradation rates must be <  $5 \times 10^{-4}$  %/hour (<500 ppm %/hour)
- Capacity loss in water-based organic systems arises from two major processes:
  1. Chemical transformations
  2. Molecular crossover



**Michael Reaction (1,4-addition)** with a nucleophile such as *water, hydroxide ion or phenoxide ion*.




- Loss of capacity
- Decrease of cell voltage

**Desulfonation** or loss of the sulfonic acid group

- Loss of solubility
- Decrease of cell voltage


# Michael Reaction or 1,4-Addition

- Occurs more readily on electron-deficient systems




# Desulfonation Processes (acid medium)

Proto-desulfonation in Acid: Loss of sulfonic acid group




# Desulfonation Processes (alkaline medium)

- Oxidative Hydroxy-desulfonation



- Nucleophilic ( $S_NAR$ ) Hydroxy-desulfonation



# Technical Approach

- To reduce the propensity for *Desulfonation* and *Michael Reactions*
  - Examine the effect of substitution by alkyl and phenyl groups on benzoquinone rings
  - Examine these reactivities in bi-functional molecules
- Adding functional groups to increase solubility without increasing reactivity.
- Developing procedures for in-house synthesis of compounds
- Follow the stability changes by NMR and GC-MS to determine effects of long-term cycling
- Electrochemical kinetics testing using RDE and CV on glassy carbon/graphite
- Establishing solubility in charged and discharged state
- Establish decay rates by extended cycling in symmetric cell configuration in flow cells (25 cm<sup>2</sup>)



# Tasks to Address Challenges


- Task 1. Synthesis, purification and scale-up of materials.
- Task 2. Electrochemical characterization of charge/discharge reversibility and electrode potential.
- Task 3. Characterization of solubility and diffusion coefficient
- Task 4. Crossover rate studies
- Task 5. Passive and active durability studies using flow cell and electrolysis
- Task 6. Reporting, Reviews and Publication

## Accomplishments in the past year


- Several promising benzoquinone derivatives have been synthesized and tested in acid and alkaline media.
- Preliminary results show these molecules possess distinct reactivity and stability based on the substituent groups.
- Verified that anthraquinone-based molecules can be stabilized to make positive side materials and also allow for bifunctional activity to achieve high cell voltage.
- Verified the long-term cycling behavior of stabilized redox materials.

# Three Classes of Benzoquinone-derived Positive Side Materials


## Alkyl-substituted




## Aryl-substituted



## Anthraquinone-derived




# Studies on 6-methyl hydroquinone-3-sulfonic acid (MMS)




MMS

## Synthesis




10 mM MMS in 1 M Sulfuric Acid; Graphite Electrode;  
50 mV/s vs MSE




- Cyclic voltammograms suggest excellent reversibility.
- Slow chemical transformations are not captured in a CV.


# Cycling studies on MMS




CV at 50 mV/s, Graphite Electrode  
in 1 M Sulfuric Acid



## NMR Analysis-Starting Material




After 516 cycles: Significant Loss  
of the Aromatic Protons and  
appearance of multiple aliphatic  
peaks




# Polymerization of MMS


- Shown: dimerization




- Process can continue (chain propagation) to form redox active oligomers





# CV Characterization of O3MMS, O4MMS, O3MDS





O3MMS, 1M H<sub>2</sub>SO<sub>4</sub>, GC, 50mV/second



O3MDS, 1M H<sub>2</sub>SO<sub>4</sub>, GC, 50mV/second




O4MMS, 1M H<sub>2</sub>SO<sub>4</sub>, GC, 50mV/second




# Tertiary-Butyl Substituted Orthobenzoquinone monosulfonic acid (O4TBMS)

## Synthesis





O4TBMS, 1M  $\text{H}_2\text{SO}_4$ , GC, 50mV/second



Potential shift to positive values upon sulfonation

1M O4TBMS/ 1M AQDS cell cycled at 100 mA/cm<sup>2</sup> in sulfuric acid




NMR Analysis confirms fast single step hydroxylation





# Aryl-Substituted benzoquinones


## PHQS, PHQDS, Aryl sulfones






PHQS, 1M H<sub>2</sub>SO<sub>4</sub>, GC, 50mV/second





PHQDS, 1M H<sub>2</sub>SO<sub>4</sub>, GC, 50mV/second



### Aryl sulfones




### Cycling of PHQS /AQDS cell in 1M sulfuric acid at 100 mA/cm<sup>2</sup>





— charge cap  
— discha cap



# Bifunctional Anthraquinone –Derived Molecules- Alizarin Red in Alkaline Media





Alizarin Red, 1M KOH, GC, 50mV/second




Charge  
Discharge



Michael Reaction




KOH





# Hydroxydesulfonation of Alizarin Red

## NMR Analysis of cycled Alizarin Red







Confirms the formation of the tetrahydroxylated product


# Purpurin-a viable pathway to using anthraquinone derivatives



Hydroxylated  
Purpurin can be  
cycled in alkali  
without noticeable  
degradation

# Other Promising Bifunctional Molecules Studied

Compound	Structure	$E_{1/2}$ (Volts) vs MMO	Bi-functional Material Potential
1,2,4-trihydroxy anthraquinone		-0.84 V	Yes, 840 mV difference
1,4-dihydroxy anthraquinone		-0.63 V	Yes, 880 mV difference
1,2-dihydroxy anthraquinone		-0.73 V	Yes, 1 V difference
1,2-dihydroxy, 3-sulfonic acid anthraquinone		-0.75 V	Yes, 930 mV difference



Potential to increase cell voltage by substitution

# Next Steps



- Increase the solubility of stable redox molecules to ensure high concentrations.
- Develop methods of sulfonation for the non-participating ring.
- Complete the characterization in alkaline media
- Down-select molecules for full-cell testing
- Pursue further molecular designs to avoid oligomer formation.
- Explore the bifunctional nature of stabilized redox molecules in full cell.

# Acknowledgements

- Dr. Imre Gyuk , DoE's Office of Electricity.
- Drs. Wei Wang, David Reed and Vincent Sprenkle at PNNL
- University of Southern California, Loker Hydrocarbon Research Institute for post-doctoral fellow and graduate student support.




# The USC Organic Flow Battery Team



Dr. Lena Hooper  
(past student)



Sri Narayan



Dr. Bo Yang



Vinayak  
Krishnamurti

Dr. Sankarganesh  
Krishnamoorthy  
(past student)



Advait Murali

Dr. Archith  
Nirmalchandar  
(past student)



Prof. Surya  
Prakash

Dr. Robert Aniszfeld

